

For Research Use Only

CoraLite® Plus 555-conjugated PLK1 Polyclonal antibody



Catalog Number: CL555-10305

Featured Product

Basic Information

Catalog Number:

CL555-10305

Size:

1 mg/mL

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0284

GenBank Accession Number:

BC002369

GeneID (NCBI):

5347

UNIPROT ID:

P53350

Full Name:

polo-like kinase 1 (Drosophila)

Calculated MW:

68 kDa

Observed MW:

62 kDa

Purification Method:

Antigen affinity purification

Excitation/Emission maxima wavelengths:

554 nm / 570 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Background Information

PLK1(Polo-like kinase 1), also named as PLK and STPK13, belongs to the protein kinase superfamily, Ser/Thr protein kinase family and CDC5/Polo subfamily. PLK1 is a Serine/threonine-protein kinase that performs several important functions throughout M phase of the cell cycle, including the regulation of centrosome maturation and spindle assembly, the removal of cohesins from chromosome arms, the inactivation of APC/C inhibitors, and the regulation of mitotic exit and cytokinesis. The localization of Polo changes during the cell cycle. In interphase, Polo shows a mostly diffuse cytoplasmic localization. Polo becomes concentrated on centrosomes from early prophase and appears on centromeres or kinetochores from late prophase, before nuclear envelope breakdown (NEB). After anaphase onset, Polo relocates to the central spindle and remains enriched at the midbody ring in late stages of cytokinesis (PMID: 29167465).

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 A549 cells were intracellularly stained with 0.2 μ g CoraLite® Plus 555 Anti-Human PLK1 (CL555-10305) (red), or 0.2 μ g CL555 Rabbit Isotype Control Antibody (CL555-30000) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).