For Research Use Only

CoraLite®555-conjugated IRF3 Polyclonal antibody

Catalog Number: CL555-11312 Featured Product



Basic Information

Catalog Number: CL555-11312 Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG1858

GenBank Accession Number:

BC009395 GeneID (NCBI): 3661 **UNIPROT ID:** Q14653 Full Name:

interferon regulatory factor 3

Calculated MW: 47 kDa Observed MW: 50-60 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Applications

Tested Applications: FC (Intra)

Species Specificity:

human

Positive Controls:

FC (Intra): human peripheral blood lymphocytes,

Background Information

The virul-induced expression of interferon(IFN) genes in infected cells implicate in the interplay of several constitutively expressed and virus-activated transcription factors. A family of IFN regulatory factors(IRFs) have been shown to has a role in the transcription of IFN genes as well as IFN-stimulated genes. IRF3 is a novel key transcriptional regulator of type I IFN-dependent immune responses and involves in the innate immune response against DNA and RNA viruses, by binding to the promoters of IFN. It located in the cytoplasm of uninfected cells in an inactive form, and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, could be phosphorylated by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization, nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and ISG genes. IRF3 exists some isoforms with MV 47-49 kDa, 33 kDa and 12-16 kDa.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1x10^6 human peripheral blood lymphocytes were intracellularly stained with 0.8 ug CoraLite®555-conjugated IRF3 Polyclonal antibody (CL555-11312)(red), or 0.8 ug CoraLite®555-conjugated Rabbit IgG control Rabbit PolyAb (CL555-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.