For Research Use Only

CoraLite®555-conjugated TIA1 Polyclonal antibody



Purification Method:

IF/ICC 1:300-1:1200

wavelengths: 557 nm / 570nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL555-12133

Featured Product

Basic Information

Catalog Number: CL555-12133

1000 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG2778

40 kDa

GenBank Accession Number:

BC015944 GeneID (NCBI): 7072 **UNIPROT ID:** P31483 Full Name:

TIA1 cytotoxic granule-associated RNA binding protein

Calculated MW: 214 aa, 24 kDa, 43 kDa Observed MW:

Positive Controls:

IF/ICC: HepG2 cells,

Applications

Tested Applications:

Species Specificity: human, mouse, rat

Background Information

TIA1, also named as p40-TIA-1, is involved in alternative pre-RNA splicing and regulation of mRNA translation by binding to AU-rich elements (AREs) located in mRNA 3' untranslated regions (3' UTRs). It possesses nucleolytic activity against cytotoxic lymphocyte target cells. TIA1 may be involved in apoptosis. Two isoforms of this protein exist - 41kDa and 42kDa. one of these was a missense variant (P362L) in TIA1. Similar to the ALS-related disease proteins TDP-43, hnRNPA1, and FUS, TIA1 is an RNA-binding protein containing a prionlike LCD and assembles into membrane-less organelles, including SGs. Postmortem neuropathology of five TIA1mutations carriers showed a consistent pathological signature with numerous round, hyaline, TAR DNA-binding protein 43 (TDP-43)-positive inclusions.TIA1mutations significantly increased the propensity of TIA1 protein to undergo phase transition. In live cells,TIA1mutations delayed stress granule (SG) disassembly and promoted the accumulation of non-dynamic SGs that harbored TDP-43. Moreover, TDP-43 in SGs became less mobile and insoluble.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite®555 TIA1 antibody (CL555-12133) at dilution of 1:600.