For Research Use Only

## CoraLite®555-conjugated VDAC1/Porin Polyclonal antibody

Catalog Number: CL555-55259

Featured Product



**Basic Information** 

Catalog Number: GenBank Accession Number: CL555-55259 NM\_003374

Concentration: GeneID (NCBI): 7416

Source: UNIPROT ID: Rabbit P21796

Isotype: Full Name:

voltage-dependent anion channel 1

Calculated MW: 31 kDa Observed MW: 31 kDa Purification Method: Antigen affinity purification Recommended Dilutions: IF-P 1:50-1:500

Excitation/Emission maxima wavelengths:

IF-Fro 1:50-1:500

557 nm / 570 nm

Applications

Tested Applications: IF-P, IF-Fro, FC (Intra) Species Specificity: human, mouse, rat Positive Controls:

IF-P: human liver cancer tissue,
IF-Fro: mouse liver tissue,

## **Background Information**

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1. C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This antibody is specific to VDAC1.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using CoraLite®555 VDAC1/Porin antibody (CL555-55259) at dilution of 1:200.



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite®555-conjugated VDAC 1/Porin Polyclonal antibody (CL555-55259) (red), or 0.8 ug CoraLite®555-conjugated Rabbit 1gG control Rabbit PolyAb (CL555-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse liver tissue using CoraLite®555 VDAC1/Porin antibody (CL555-55259) at dilution of 1:200.