

For Research Use Only

CoraLite®555-conjugated PPAR Gamma Monoclonal antibody



Catalog Number:CL555-60127

Basic Information

Catalog Number: CL555-60127	GenBank Accession Number: BC006811	Purification Method: Protein G purification
Size: 1000 µg/ml	GeneID (NCBI): 5468	CloneNo.: 4E12F10
Source: Mouse	UNIPROT ID: P37231	Recommended Dilutions: IF 1:50-1:500
Isotype: IgG1	Full Name: peroxisome proliferator-activated receptor gamma	Excitation/Emission maxima wavelengths: 557 nm / 570nm
Immunogen Catalog Number: AG10005	Calculated MW: 58 kDa	
	Observed MW: 50-60 kDa	

Applications

Tested Applications: IF-P	Positive Controls: IF : human colon tissue,
Species Specificity: human	

Background Information

Peroxisome Proliferator-Activated Receptors (PPARs) are ligand-activated intracellular transcription factors, members of the nuclear hormone receptor superfamily (NR), that includes estrogen, thyroid hormone receptors, retinoic acid, Vitamin D3 as well as retinoid X receptors (RXRs). The PPAR subfamily consists of three subtypes encoded by distinct genes denoted PPAR α (NR1C1), PPAR β / δ (NR1C2) and PPAR γ (NR1C3), which are activated by selective ligands. PPAR γ , also named as PPARG, contains one nuclear receptor DNA-binding domain and is a receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. It plays an important role in the regulation of lipid homeostasis, adipogenesis, INS resistance, and development of various organs. Defects in PPARG are the cause of familial partial lipodystrophy type 3 (FPLD3) and may be associated with susceptibility to obesity. Defects in PPARG can lead to type 2 INS-resistant diabetes and hypertension. PPARG mutations may be associated with colon cancer. Genetic variations in PPARG are associated with susceptibility to glioma type 1 (GLM1). PPARG has two isoforms with molecular weight 57 kDa and 54 kDa (PMID: 9831621), but modified PPARG is about 67 kDa (PMID: 16809887). PPARG2 is a splice variant and has an additional 30 amino acids at the N-terminus (PMID: 15689403). Experimental data indicate that a 45 kDa protein displaying three different sequences immunologically related to the nuclear receptor PPARG2 is located in mitochondria (mt-PPAR). However, the molecular weight of this protein is clearly less when compared to that of PPARG2 (57 kDa). (PMID: 10922459). PPARG has been reported to be localized mainly (but not always) in the nucleus. PPARG can also be detected in the cytoplasm and was reported to possess extra-nuclear/non-genomic actions (PMID: 17611413; 19432669; 14681322).


Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human colon tissue using CoraLite®555 PPAR Gamma antibody (CL555-60127, Clone: 4E12F10) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed human colon tissue using CoraLite®555 PPAR Gamma antibody (CL555-60127, Clone: 4E12F10) at dilution of 1:200.