For Research Use Only

## CoraLite®555-conjugated Ecadherin Monoclonal antibody



**Purification Method:** 

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:50-1:500

wavelengths: 557 nm / 570 nm

CloneNo.:

6B11F11

Catalog Number: CL555-60335

**Featured Product** 

**Basic Information** 

Catalog Number: CL555-60335

1000 µg/ml Source: Mouse

Size:

Isotype:

lgG2b Immunogen Catalog Number:

AG15085

**Tested Applications:** 

Species Specificity: human, pig

GenBank Accession Number:

BC141838 GeneID (NCBI):

**UNIPROT ID:** P12830 Full Name:

(epithelial)

120 kDa

cadherin 1, type 1, E-cadherin

Calculated MW: 882 aa, 97 kDa Observed MW:

Positive Controls:

IF/ICC: MCF-7 cells,

## **Background Information**

**Applications** 

Cadherins are a family of transmembrane glycoproteins that mediate calcium-dependent cell-cell adhesion and play an important role in the maintenance of normal tissue architecture. E-cadherin (epithelial cadherin), also known as CDH1 (cadherin 1) or CAM 120/80, is a classical member of the cadherin superfamily which also include N-, P-, R-, and B-cadherins. E-cadherin is expressed on the cell surface in most epithelial tissues. The extracellular region of E-cadherin establishes calcium-dependent homophilic trans binding, providing specific interaction with adjacent cells, while the cytoplasmic domain is connected to the actin cytoskeleton through the interaction with p120-,  $\alpha$  -,  $\beta$  -, and  $\gamma$  -catenin (plakoglobin). E-cadherin is important in the maintenance of the epithelial integrity, and is involved in mechanisms regulating proliferation, differentiation, and survival of epithelial cell. E-cadherin may also play a role in tumorigenesis. It is considered to be an invasion suppressor protein and its loss is an indicator of high tumor aggressiveness.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using Coralite®555 E-cadherin antibody (CL555-60335, Clone: 6B11F11) at dilution of 1:200.