For Research Use Only

CoraLite® Plus 555 Anti-Human CD7 (4H9)

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: CL555-65203

Basic Information

Catalog Number: GenBank Accession Number: CL555-65203 BC009293
Concentration: GeneID (NCBI):

100tests, 5 ul/test 9

Source: ENSEMBL Gene ID: Mouse ENSG00000173762
Isotype: UNIPROT ID:

IgG2a, kappa P09564
Full Name:
CD7 molecule

Calculated MW: 240 aa, 25 kDa

Purification Method:

Affinity purification

CloneNo.: 4H9

Recommended Dilutions: FC: 5 ul per 10^6 cells in 100 $\,\mu$ l

suspension

Excitation/Emission maxima

wavelengths: 554 nm / 570 nm

Applications

Tested Applications:

FC

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

Background Information

CD7, also known as Leu-9 or GP40, is a 40-kDa single-pass type I transmembrane glycoprotein belonging to the immunoglobulin superfamily. CD7 is expressed on thymocytes, T cells, NK cells, and cells in the early stages of T, B, and myeloid cell differentiation (PMID: 10530432). It is one of the earliest antigens to appear on cells of the T-lymphocyte lineage (PMID: 3501369). CD7 plays a significant role in T-cell and T-cell/B-cell interactions during early lymphoid development. It is involved in the regulation of cell signaling and biology, including T-cell activation, proliferation, and the expression of interleukin-2 receptor alpha (IL-2R a) (PMID: 11485208). CD7 is not only a marker for T-cells but also has functional implications in immune cell interactions and adhesion. It can provide co-stimulatory signals and is associated with CD3 and CD45, participating in T-cell activation (PMID: 7523512).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3 $\,$

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with APC Anti-Human CD4 and 5 ul CoraLite® Plus 555 Anti-Human CD7 (CL555-65203, Clone:4H9) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.