## For Research Use Only

## CoraLite®555 Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody



Catalog Number: CL555-65559

**Basic Information** 

Catalog Number: CL555-65559 Concentration:

100tests, 5 ul/test Source: Mouse Isotype:

IgG2a

ENSEMBL Gene ID: ENSG00000153563 UNIPROT ID: P01732 Full Name: CD8a molecule

> Calculated MW: 235 aa, 26 kDa

BC025715

GeneID (NCBI):

GenBank Accession Number:

**Purification Method:** 

Protein A purification CloneNo.:

UCHT4

Recommended Dilutions: FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

## **Background Information**


 ${\tt CD8}\ is\ a\ transmembrane\ glycoprotein\ composed\ of\ two\ disulfide-linked\ chains.\ It\ can\ be\ present\ as\ a\ homodimer\ of\ present\ constant and\ constant\ constant$ CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with APC Anti-Human CD3, and 5 ul CoraLite®555 Anti-Human CD8 (UCHT4) Mouse IgG2a RecAb (CL555-65559, Clone:UCHT4) or CoraLite® Plus 555 Mouse IgG2a Isotype Control (C1.18.4) (CL555-65208, Clone: C1.18.4). Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite®555 Anti-Human CD8 (UCHT4) Mouse IgG2a RecAb (CL555-65559, Clone: UCHT4) (red) or CoraLite® Plus 555 Mouse IgG2a Isotype Control (C1.18.4) (CL555-65208, Clone: C1.18.4) (blue). Cells were not fixed.