For Research Use Only

CoraLite®555-conjugated Phospho-Hamartin/TSC1 (Ser511) Recombinant antibody




Catalog Number: CL 555-80340

Basic Information	Catalog Number: CL555-80340	GenBank Accession Number: NM_000368	Purification Method: Protein A purification
	Size: 1000 µg/ml	GenelD (NCBI): 7248	CloneNo.: 3N6
	Source: Rabbit Isotype: IgG	UNIPROT ID: Q92574 Full Name: tuberous sclerosis 1	Excitation/Emission maxima wavelengths: 557 nm / 570 nm
		Observed MW: 150 kDa	
Applications	Tested Applications: FC (Intra)		
	Species Specificity: human, mouse		
Background Information	TSC1, also named as Hamartin, is a tumor suppressor gene syndrome whose manifestations can include seizures, mental retardation, autism, and tumors in the brain, retina, kidney, heart, and skin. The tuberous sclerosis 1 (TSC1)/TSC2 tumor suppressor complex serves as a repressor of the mTOR pathway, and disruption of TSC1/TSC2 complex function may contribute to tumorigenesis. TSC1 is phosphorylated by IKKb at Ser487 and Ser511 in vitro and in vivo, and phosphorylation at these sites enhances dissociation of the TSC1/TSC2 complex, which in turn induces mTOR activation. (PMID: 17693255). TSC1 is also threonine-phosphorylated at three sites (Thr417, Ser584, and Thr1047) in a reaction catalyzed by cyclin-dependent kinase 1 (CDK1) (PMID: 14551205).		
Storage	Storage: Store at -20°C. Avoid exposure to lig Storage Buffer: PBS with 50% Glycerol, 0.05% Procl	ht. Stable for one year after shipment.	
	Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 HeLa cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug CoraLite®555 Anti-Human Phospho-Hamartin/TSC1 (Ser511) (CL555-80340, Clone:3N6) (red), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with 50% MeOH.