For Research Use Only

CoraLite®555-conjugated IBA1 Recombinant antibody

Catalog Number: CL555-81728




Basic Information	Catalog Number: CL555-81728	GenBank Accession Number: BC009474	Purification Method: Protein A purification
	Size: 1000 µg/ml	GenelD (NCBI): 199	CloneNo.: 4D5
	Source: Rabbit	ENSEMBL Gene ID: ENSG00000204472	Recommended Dilutions: IF-P 1:50-1:500
	Isotype: IgG Immunogen Catalog Number: AG1363	UNIPROT ID: P55008 Full Name: allograft inflammatory factor 1	Excitation/Emission maxima wavelengths: 557 nm / 570 nm
		Observed MW: 17 kDa	
		Applications	
Species Specificity: Human, mouse, rat			
Background Information	IBA1 is a 143 amino acid cytoplasmic, inflammation response scaffold protein. It is constitutively expressed in monocytes and macrophages and is known to be involved in macrophage activation. It is a marker of activated macrophage. Expression of IBA1 is up-regulated in activated microglia following facial nerve axotomy, ischemia, and several brain diseases, thereby implicating it in the activated phenotypes of microglia.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using Coralite®555 IBA1 antibody (CL555-81728, Clone: 4D5) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using Coralite®555 IBA1 antibody (CL555-81728, Clone: 4D5) at dilution of 1:200.



1X10^6 THP-1 cells were intracellularly stained with 0.8 ug CoraLite®555 Anti-Human IBA1 (CL555-81728, Clone:4D5) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).