For Research Use Only

CoraLite®594-conjugated CUL4B Polyclonal antibody

Catalog Number:CL594-12916 Featured Product



Basic Information

Catalog Number: CL594-12916 Concentration: 1000 ug/ml

Source:
Rabbit
Isotype:
IgG
Immunogen Catalog Number:

AG3563 913 aa, 104 kDa Observed MW: 102 kDa Purification Method: Antigen affinity purification Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths: 588 nm / 604 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse Positive Controls:

IF/ICC: HepG2 cells, HeLa cells

Background Information

Cullin-RING ligases (CRLs) complexes participate in the regulation of diverse cellular processes, including cell cycle progression, transcription, signal transduction and development (PMID: 21816341)(PMID: 21554755). Serving as the scaffold protein, cullins are crucial for the assembly of ligase complexes, which recognize and target various substrates for proteosomal degradation. Two cullin 4 (CUL4) proteins, CUL4A (87 kDa) and CUL4B(104 kDa), are two members in cullin family with 83% of identity. Mutations in human CUL4B are one of the major causes of X-linked mental retardation. Cul4b knockout mice demonstrated that CUL4B is indispensable for embryonic development in the mouse (PMID: 22606329).

GenBank Accession Number:

BC036216

8450

GeneID (NCBI):

UNIPROT ID:

Full Name:

cullin 4B
Calculated MW:

Q13620


Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA Aliquoting is unnecessary for -20° C storage


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®594 CUL4B antibody (CL594-12916) at dilution of 1:200, CL488-phalloidin (green).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® 594 CUL4B antibody (CL594-12916) at dilution of 1:200, CL488-phalloidin (green).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human CUL4B (CL594-12916) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).