For Research Use Only

CoraLite®594-conjugated LC3 Polyclonal antibody


Catalog Number:CL594-14600 Featured Product




Basic Information	Catalog Number: CL594-14600	GenBank Accession Number: BC067797	Purification Method: Antigen affinity purification	
	Size: 1000 µg/ml	GenelD (NCBI): 81631	Recommended Dilutions: IF/ICC 1:50-1:500	
	Source: Rabbit Isotype: IgG	ENSEMBL Gene ID: ENSG00000140941 UNIPROT ID: Q9GZQ8	Excitation/Emission maxima wavelengths: 588 nm / 604 nm	
				Immunogen Catalog Number: AG6144
	Calculated MW: 15 kDa			
	Observed MW: 14-18 kDa			
	Applications	Tested Applications: IF/ICC, FC (Intra)	Positive Controls:	
Species Specificity: human, mouse, rat		es Specificity:		
Background Information	Map1LC3, also known as LC3, is the human homolog of yeast Atg8 and is involved in the formation of autophagosomal vacuoles, called autophagosomes. Three human Map1LC3 isoforms, MAP1LC3A, MAP1LC3B, and MAP1LC3C, undergo post-translational modifications during autophagy. And they differ in their post-translation modifications during autophagy. Map1LC3 also exists in two modified forms, an 18 kDa cytoplasmic form that was originally identified as a subunit of the microtubule-associated protein 1, and a 14-16 kDa form that is associated with the autophagosome membrane. This antibody can cross react with MAP1LC3A, MAP1LC3B, and MAP1LC3C.			
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage			

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Chloroquine treated HepG2 cells using CoraLite®594 LC3 antibody (CL594-14600) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human LC3 (CL594-14600) (red), or 0.4 ug CoraLite®594-conjugated Rabbit 1gG control Rabbit PolyAb (CL594-30000, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).