For Research Use Only


CoraLite®594-conjugated NRF2, NFE2L2 Polyclonal antibody Catalog Number:CL594-16396 Featured Product 1 Publications

oroteintec Antibodies | ELISA kits | Proteins www.ptglab.com


| Basic Information | Catalog Number: CL594-16396 | GenBank Accession Numl BC011558 | per: Purification Method: Antigen affinity purification | ation | |
|------------------------|---|---|--|--|--|
| | Size: 1000 ug/ml | GenelD (NCBI): 4780 | Recommended Dilution IF/ICC 1:50-1:500 | IS: | |
| | Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG9489 | UNIPROT ID: Q16236 | Excitation/Emission ma wavelengths: | Excitation/Emission maxima wavelengths: | |
| | | Full Name: 588 nm / 604 nm nuclear factor (erythroid-derived 2)- like 2 Calculated MW: | | | |
| | | 605 aa, 68 kDa Observed MW: 110 kDa | | | |
| Applications | Tested Applications: IF/ICC | Positive Controls: IF/ICC : HepG2 cells, | | | |
| | Cited Applications: | | | | |
| | Species Specificity: human | | | | |
| Background Information | NRF2, also named as NFE2L2, belongs to the bZIP family and CNC subfamily. It is a transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. NRF2 is important for the coordinated up-regulation of genes in response to oxidative stress. It may be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region. Nrf2 is a key player in the regulation of genes encoding for many antioxidative response enzymes. The expression of NRF2 may be induced under oxidative stress (PMID:14567983).In lung cancer, Nrf2 activation in malignant cells has been associated with tumor progression and chemotherapy resistance(PMID:20534738). Identifying patients with abnormal NRF2 expression may be important for selection for chemotherapy in NSCLC. As new investigators break into the emerging field of Nrf2 research, confusion regarding the correct migratory pattern of Nrf2 is causing doubts about the accuracy and reproducibility of published results. This letter provides solid evidence that the actually observed molecular weight of Nrf2 is about 70kDa and 95-110 kDa. (PMID: 22703241). | | | | |
| Notable Publications | Author | Pubmed ID Journal | Ap | plication | |
| | Qi Li | 36974993 Am J Chi | n Med IF | | |
| Storage | Storage: Store at -20°C. Avoid exposure to Storage Buffer: PBS with 50% Clycerol, 0.05% Pr Aliquoting is unnecessary for -20 | oclin300, 0.5% BSA, pH 7.3. | shipment. | | |

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594 NRF2, NFE2L2 antibody (CL594-16396) at dilution of 1:100.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using Coralite®594 NRF2, NFE2L2 antibody (CL594-16396) at dilution of 1:200.