For Research Use Only

CoraLite®594-conjugated VEGFA Polyclonal antibody



Purification Method:

IF/ICC 1:50-1:500

wavelengths: 588 nm / 604 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL594-19003

Featured Product

Basic Information

Catalog Number: CL594-19003

Size: 1000 µg/ml Source: Rabbit Isotype:

IgG vascular endothelial growth factor A
Immunogen Catalog Number: Calculated MW:

AG13500 412 aa, 46 kDa

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse Positive Controls: IF/ICC : HeLa cells,

Background Information

VEGFA, also named as VEGF or VPF, belongs to the PDGF/VEGF growth factor family. It is a growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. VEGFA induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. Defects in VEGFA are associated with microvascular complications of diabetes type 1 (MVCD1). VEGFA has 17 isoforms with MW from 16 to 45 kDa. Some isoforms have homodimer forms (e.g.; VEGFA189 38kDa or VEFGA110 34kDa). VEGF-A exists in at least seven homodimeric isoforms. The monomers consist of 121, 145, 148, 165, 183, 189, or 206 amino acids (PMID:15602010). This antibody can recognize all VEGFA isoforms.

GenBank Accession Number:

BC065522

7422

P15692

GeneID (NCBI):

UNIPROT ID:

Full Name:

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite®594 VEGF antibody (CL594-19003) at dilution of 1:100.



1X10^6 NIH/3T3 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human VEGF (CL594-19003) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).