## For Research Use Only

## CoraLite®594-conjugated Ki-67 Polyclonal antibody

Catalog Number:CL594-27309

Featured Product

1 Publications



**Basic Information** 

Catalog Number: CL594-27309

Size: 1000 ug/ml Source: Rabbit

Immunogen Catalog Number:

AG26266

Isotype:

GenBank Accession Number:

NM\_002417 GeneID (NCBI): 4288

Full Name: antigen identified by monoclonal antibody Ki-67

Calculated MW: 359 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra) Cited Applications:

IF

Species Specificity:

human **Cited Species:** human

Positive Controls: IF/ICC: HeLa cells,

**Background Information** 

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Liang Xu	39054954	Oncol Rep	IF

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite® 594 Kl67 antibody (CL594-27309) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®594 Kl67 antibody (CL594-27309) at dilution of 1:200.



1x10^6 Ramos cells were intracellularly stained with 0.8 ug CoraLite® 594 Anti-Human Kl67 (CL594-27309)(red), or 0.8 ug CoraLite® 594-conjugated Rabbit IgG control Rabbit PolyAb (CL594-30000) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.