For Research Use Only

## CoraLite®594-conjugated Follistatin Monoclonal antibody



Catalog Number: CL594-60060

**Basic Information** 

Catalog Number: GenBank Accession Number: CL594-60060 BC004107

CL594-60060 BC004107
Size: GeneI D (NCBI):

 $\begin{array}{lll} 1000 \ \mu \ g/ml & 10468 \\ \\ Source: & UNIPROT \ ID: \\ \\ Mouse & P19883 \\ \\ Isotype: & Full \ Name: \end{array}$ 

IgG2b follistatin
Immunogen Catalog Number: Calculated MW:

AG0775 33 aa, 7 kDa

Observed MW: 35 kDa Purification Method:

Caprylic Acid-Ammonium Sulfate

Precipitation
CloneNo.:
4E9B10

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths:

588 nm / 604 nm

**Applications** 

**Tested Applications:** 

IF/ICC

Species Specificity: human, mouse

Positive Controls:

IF/ICC: HeLa cells,

**Background Information** 

Follistatin (FST) is a member of the tissue growth factor  $\beta$  family and is a secreted glycoprotein that antagonizes many members of the family, including activin A, growth differentiation factor11, and myostatin. It binds activin A with high affinity and whose expression can be induced by activin A and several other pro-inflammatory cytokines. Activin A-follistatin complexes are biologically inactive and bind to cell surface heparan sulphate-containing proteoglycans for internalisation and degradation. It has also been found to play a significant role in the management of skeletal muscle size and mass. It also has important roles in early embryonic development, differentiation of ovarian granulosa cells, liver fibrosis and polycystic ovarian syndrome.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using CoraLite®594 Follistatin antibody (CL594-60060, Clone: 4E9B10) at dilution of 1:200.