

For Research Use Only

CoraLite®594-conjugated Chromogranin A Monoclonal antibody

Catalog Number:CL594-60135



Basic Information

Catalog Number:

CL594-60135

Concentration:

1000 ug/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG0807

GenBank Accession Number:

BC006459

GeneID (NCBI):

1113

UNIPROT ID:

P10645

Full Name:

chromogranin A (parathyroid
secretory protein 1)

Calculated MW:

51 kDa

Observed MW:

70 kDa

Purification Method:

Protein G purification

CloneNo.:

3H11C11

Recommended Dilutions:

IF/ICC 1:50-1:500

Excitation/Emission maxima
wavelengths:

588 nm / 604 nm

Applications

Tested Applications:

IF/ICC, FC (Intra)

Species Specificity:

human, mouse, rat, pig

Positive Controls:

IF/ICC : Neuro-2a cells,

Background Information

Chromogranin A is a member of the granin family of neuroendocrine secretory proteins. It is located in secretory vesicles of neurons and endocrine cells. Chromogranin A is the precursor to several functional peptides including vasostatin, pancreastatin, catestatin and parastatin. These peptides negatively modulate the neuroendocrine function of the releasing cell (autocrine) or nearby cells (paracrine). CgA is one of the most used tumor markers in NET's (neuroendocrine tumors), and elevated CgA concentrations have been demonstrated in serum or plasma of patients with different types of these tumors.

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1×10^6 SH-SY5Y cells were intracellularly stained with 0.8 μ g CoraLite® 594 Anti-Human Chromogranin A (CL594-60135, Clone:3H11C11) (red), or 0.8 μ g Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1×10^6 Neuro-2a cells were intracellularly stained with 0.8 μ g CoraLite® 594 Anti-Human Chromogranin A (CL594-60135, Clone:3H11C11) (red), or 0.8 μ g Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Ethanol) fixed Neuro-2a cells using CoraLite® 594 Chromogranin A antibody (CL594-60135, Clone: 3H11C11) at dilution of 1:200.