For Research Use Only

CoraLite® Plus 594 Anti-Human CD68 (KP1)

Antibodies | ELISA kits | Proteins www.ptglab.com


Catalog Number: CL594-65202


Basic Information	Catalog Number: CL594-65202	GenBank Accession Number: BC015557	Purification Method: Protein G purification				
	Size: 100tests, 5 ul/test	GenelD (NCBI): 968	CloneNo.: KP1				
	Source: Mouse Isotype: IgG1, kappa	UNIPROT ID: P34810 Full Name: CD68 molecule Calculated MW: 37 kDa	Excitation/Emission maxima wavelengths: 594 nm / 615 nm				
				Applications	Tested Applications: FC (Intra)		
					Species Specificity: human		
Background Information	CD68 is a type I transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It belongs to the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family and primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. CD68 is also a member of the scavenger receptor family. It may play a role in phagocytic activities of tissue macrophages.						
Storage	Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.1% sodium azide and 0.5% BSA, pH 7.3.						

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 human PBMCs were intracellularly stained with 5 ul CoraLite®594 Anti-Human CD68 (CL594-65202, Clone:KP1) or CoraLite®594 Mouse IgG1 Isotype Control (CL594-65124, Clone: MOPC-21). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011) prior to staining. Monocytes were gated.

1x10^6 human PBMCs were intracellularly stained with 5 ul CoraLite®594 Anti-Human CD68 (CL594-65202, Clone:KP1) or CoraLite®594 Mouse IgG1 Isotype Control (CL594-65124, Clone: MOPC-21). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011) prior to staining. Monocytes were gated.