For Research Use Only

CoraLite®594-conjugated HDAC1 Monoclonal antibody

Catalog Number: CL594-66085




Basic Information	Catalog Number: CL594-66085	GenBank Accession Number: BC000301	Purification Method: Protein G purification				
	Size: 1000 ug/ml	GeneID (NCBI): 3065	CloneNo.: 4C4G7				
	Source: Mouse	UNIPROT ID: Q13547	Recommended Dilutions: IF/ICC 1:4000-1:16000				
	Isotype: IgG1 Immunogen Catalog Number: AG19128	Full Name: histone deacetylase 1 Calculated MW: 55 kDa Observed MW: 60-65 kDa	Excitation/Emission maxima wavelengths: 588 nm / 604 nm				
				Applications	Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : A431 cells, HeLa cells	
					Species Specificity: human, mouse, rat		
Background Information	Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex, which is responsible for gene expression silencing. It also plays an important role in the control of cell proliferation and differentiation by interacting with RB, p53 and other transcription factors. At least 4 classes of HDAC were identified As a class I HDAC, HDAC 1 was primarily found in the nucleus. This antibody is raised against residues near the C terminus of human HDAC1. The calculated molecular weight of HDAC1 is 55 kDa, but modified HDAC1 is about 60-65 kDa.						
	terminus of human HDAC 1. The ca						


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CL594-66085 (HDAC1 antibody) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite®594 Anti-Human HDAC1 (CL594-66085, Clone:4C4G7) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using Coralite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using Coralite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.