For Research Use Only

CoraLite®594-conjugated MPO Monoclonal antibody Catalog Number:CL594-66177 Featured Product 1 Public







Basic Information	Catalog Number: CL594-66177	GenBank Accession Number: BC 130476	Purification Method: Thiophilic affinity chromatograph	
	Size: 1000 µg/ml	GeneID (NCBI): 4353	CloneNo.: 4C11F6	
	Source: Mouse	UNIPROT ID: P05164	Recommended Dilutions: IF-P 1:50-1:500	
	Isotype: IgA Immunogen Catalog Number: AG17564	Full Name: myeloperoxidase	Excitation/Emission maxima wavelengths: 588 nm / 604 nm	
		Calculated MW: 745 aa, 84 kDa		
		Observed MW: 90 kDa		
Applications	Tested Applications: IF-P, FC (Intra)	Positive Controls:		
	Cited Applications:	IF-P : human appendicitis tissue, human liver tissue		
	 Species Specificity: human, rat			
	Cited Species: human			
Background Information	The MPO gene encodes myeloperoxidase, a lysosomal hemoprotein located in the azurophilic granules of polymorphonuclear (PMN) leukocytes and monocytes. In response to stimulation, MPO is activated into a transient intermediate with potent antimicrobial oxidizing abilities(PMID:17650507). The mRNA is translated into a single protein of 90 kDa, which displays enzymatic activity and undergoes proteolytic maturation into a heavy chain of 59 kDa and a light chain of 13.5 kDa; these subunits then dimerize into the mature tetramer and the mature MPO is a heterotetramer composed of two identical heavy chains and two identical light chains(PMID:12773517). The 24-kDa material had a map identical to that of 13.5 kDa subunit and represents a dimer of the 13.5 kDa subunit (PMID:3008892). Defects in MPO are the cause of myeloperoxidase deficiency (MPOD). It has 3 isoforms produced by alternative splicing.			
Notable Publications	Author	Pubmed ID Journal	Application	
	Xing Liu	37734390 Thromb Haer		


For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using CoraLite®594 MPO antibody (CL594-66177, d Clone: 4C11F6) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using CoraLite®594 MPO antibody (CL594-66177, Clone: 4C11F6) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

1X10^6 HL-60 cells were intracellularly stained with 0.8 ug CoraLite® 594 Anti-Human MPO (CL594-66177, Clone:4C11F6) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).