For Research Use Only

CoraLite®594-conjugated BAG6 Monoclonal antibody

www.ptglab.com

Catalog Number: CL594-66661

Featured Product

Basic Information

Catalog Number: GenBank Accession Number: CL594-66661 BC003133 GeneID (NCBI): Source: Mouse 7917 **UNIPROT ID:** Isotype:

lgG1 P46379 Full Name: Immunogen Catalog Number:

AG24525 HLA-B associated transcript 3

> Calculated MW: 119 kDa Observed MW: 150-160 kDa

Purification Method: Protein G purification

CloneNo.:

1B8D3

Recommended Dilutions:

WB: 1:500-1:2000

FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications: WB, FC (Intra)

human, mouse, rat

Species Specificity:

Positive Controls:

WB: HeLa cells, LNCaP cells, HepG2 cells

FC (Intra): HeLa cells,

Background Information

BAT3 also known as Scythe or BAG6, is a nuclear protein implicated in the control of apoptosis and natural killer (NK) cell-dendritic cell (DC) interaction. BAT3 was first discovered as a member of a group of genes located within the class III region of the human major histocompatibility complex on chromosome 6, and has been extensively studied for its role in regulating apoptosis under various stress conditions such as DNA damage and endoplasmic reticulum-related stress. BAT3 has been shown to be required for p53 acetylation, which is critical for the enhancement of p53 transcriptional activity in response to DNA damage. In addition, BAT3 is involved in the regulation of development and reproduction of mammals by acting as a co-chaperone of the heat shock protein HSP70.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL594-66661 (BAG6 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human BAG6 (CL594-66661, Clone:1B8D3) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.