For Research Use Only

CoraLite®594-conjugated vwf Monoclonal antibody



Catalog Number: CL594-66682

Basic Information

Catalog Number:

CL594-66682

Size: 1000 µg/ml

Source:

Mouse

Isotype: lgG1

Immunogen Catalog Number:

AG25578

GenBank Accession Number:

GeneID (NCBI):

7450

Full Name:

von Willebrand factor

Purification Method:

Protein G purification

CloneNo.: 3F9F3

Recommended Dilutions:

IF 1:50-1:500

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

IF-P

Species Specificity:

Human

Positive Controls:

IF: human breast cancer tissue,

Background Information

 $Von \,Willebrand \,factor \,(VWF)\,is\, a\, large \,multimeric \,gly coprotein \,found \,in \,blood \,plasma \,involved \,in \,hemostasis$ following vascular injury. Due to the multimeric nature of VWF, it can range in size from 500 to 20,000 kDa due to the differences in the number of subunits comprising the protein. Each subunit is approximately 250 kDa (PMID: 9759493). The biosynthesis of VWF in vivo is limited to endothelial cells (PMID: 4209883) and megakaryocytes (PMID: 2413071). VWF synthesized in endothelial cells is either released directly into the plasma via 27186a secretory pathway, or tubulized and stored in organelles unique to this cell type called Weibel-Palade bodies (PMID: 16459301). Whereas VWF synthesized in megakaryocytes is stored in the alpha granules of platelets (PMID: 2046403). The primary function of VWF is as an adhesive plasma glycoprotein, particularly factor VIII; an essential blood-clotting protein (PMID: 6982084). VWF is also important in platelet adhesion to wound sites by binding specifically to type I and type III collagen (PMID: 11098050), with larger VWF multimers being most effective (PMID: 24448155).


Storage

Store at -20°C. Avoid exposure to light.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using CoraLite®594 vwf antibody (CL594-66682, Clone: 3F9F3) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using Coralite®594 vwf antibody (CL594-66682, Clone: 3F9F3) at dilution of 1:200.