For Research Use Only

CoraLite®594-conjugated RALYL Monoclonal antibody



Catalog Number: CL594-67534

Basic Information

Catalog Number: GenBank Accession Number:

 CL594-67534
 BC031090

 Size:
 GeneID (NCBI):

 1000 μg/ml
 138046

 Source:
 UNIPROT ID:

 Mouse
 Q86SE5

 Isotype:
 Full Name:

IgG1 RALY RNA binding protein-like

Immunogen Catalog Number:Calculated MW:AG11676291 aa, 32 kDa

Observed MW: 35-40 kDa Purification Method:

Protein G purification

CloneNo.: 1F3F4

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat, pig Positive Controls:

IF/ICC: HepG2 cells,

Background Information

RALY RNA binding protein-like (RALYL) belongs to the RALY subfamily. RALYL locates in 8q21.2, the function of which is RNA-binding and nucleotide binding. The RALYL high expression in the normal adrenal, kidney and brain (Shyamsundar et al., 2005). Furthermore, RALYL expression down-regulated correlates with mental disorder (Lee et al., 2007), Parkinson's disease (Moran et al., 2006), brain cancer (Maris et al., 2008), adrenal cancer (Giordano et al., 2009) and kidney cancer (Higgins et al., 2003). The expression of RALYL was significantly low in Clear Cell Renal Cell Carcinoma and correlated with a poor prognosis in a large number of clinical samples. Our findings showed that RALYL may be a potential therapeutic target as well as a poor prognostic

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®594 RALYL antibody (CL594-67534, Clone: 1F3F4) at dilution of 1:100.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®594 RALYL antibody (CL594-67534, Clone: 1F3F4) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human RALYL (CL594-67534, Clone:1F3F4) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).