For Research Use Only

CoraLite®594-conjugated EIF4A2 Monoclonal antibody



Purification Method:

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:50-1:500

588 nm / 604 nm

wavelengths:

CloneNo.:

2D11E11

Catalog Number: CL594-67799

Basic Information

Catalog Number: CL594-67799

Size: 1000 µg/ml Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

Tested Applications:

IF/ICC, FC (Intra)

Species Specificity: Human, Rat, Mouse

AG9085

46 kDa

GenBank Accession Number:

BC012547 GeneID (NCBI): 1974 **UNIPROT ID:** Q14240

Full Name: eukaryotic translation initiation factor 4A, isoform 2

Calculated MW: 408 aa, 46 kDa Observed MW:

Positive Controls:

IF/ICC: HEK-293 cells,

Background Information

Eukaryotic initiation factor 4A (EIF4A) has an essential role in the binding of mRNA to the 43S preinitiation complex when protein synthesis begins, as EIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon. EIF4A2 is a ATP-dependent RNA helicase, which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome.

Storage


Applications

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using Coralite®594 EIF4A2 antibody (CL594-67799, Clone: 2D11E11) at dilution of 1:200.



1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human EIF4A2 (CL594-67799, Clone:2D11E11) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).