For Research Use Only

CoraLite®594-conjugated RPA2 Monoclonal antibody

Catalog Number: CL594-67999



Basic Information

Isotype: Full Name:

IgG1 replication protein A2, 32kDa

Immunogen Catalog Number:Calculated MW:AG2478329 kDa

Observed MW: 30 kDa Purification Method: Protein G purification

CloneNo.: 1B9H11

Recommended Dilutions: IF/ICC: 1:500-1:2000

FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

IF/ICC, FC (Intra)

Species Specificity:
human, mouse, rat, pig

Positive Controls:

IF/ICC : A431 cells,

FC (Intra) : A431 cells,

Background Information

RPA2, a subunit of the heterotrimeric replication protein A (RPA) complex, is overexpressed in various cancers. RPA2 is a major target for upstream kinases, including CDKs and the three PIKKs (ATR, ataxia-telangiectasia mutated [ATM], and DNA-PK). RPA2 comprises multiple critical Ser/Thr residues that are phosphorylated in sequential order in response to genotoxic stress.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite®594 RPA2 antibody (CL594-67999, Clone: 1B9H11) at dilution of 1:1000.



1X10^6 A431 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human RPA2 (CL594-67999, Clone:1B9H11) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).