For Research Use Only

CoraLite®594-conjugated Phospho-MEK1 (Thr386) Recombinant antibody

Catalog Number:CL594-81304 Featured Product




Basic Information	Catalog Number: CL594-81304	GenBank Accession Number: BC139729	Purification Method: Protein A purification	
	Concentration: 1000 ug/ml	GenelD (NCBI): 5604	CloneNo.: 6K5	
	Source: Rabbit Isotype: IgG	ENSEMBL Gene ID: ENSG00000169032	Excitation/Emission maxima wavelengths: 588 nm / 604 nm	
		UNIPROT ID: Q02750		
		Full Name: mitogen-activated protein kinase kinase 1	nase	
		Calculated MW: 43 kDa		
		Observed MW: 40-50 kDa		
Applications	Tested Applications: FC (Intra)			
	Species Specificity: human			
Background Information	MAP2K1 encodes MAPK1, also known as MEK1. MEK1 variants can enhance MEK1 expression and ERK1 phosphorylation that together lead to continuous activation of MEK/ERK signaling pathway. MEK1 bind directly to ERK2 through a region in the N terminus of MEK. In addition, a proline-rich (PR) regulatory sequence in MEK is also involved in MEK-ERK association and signal propagation. The coupling between MEK1 and ERK2 is enhanced through phosphorylation on S298 in the MEK1 PR region, whereas phosphorylation on MEK1 T292 releases the complex. MEK1 T292 is a substrate of ERK2, but the site is also phosphorylated at a basal level when ERK2 is inhibited, suggesting several regulators of this site . Although the S298 site in MEK2 has been conserved, it lacks the T292 phosphorylation site, and it is not a substrate of PAK1. (PMID: 31972311, PMID: 17928366, PMID: 22177953)			
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3 Aliquoting is unnecessary for -20°C storage			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 HeLa cells untreated (dashed lines) or treated with λ phosphatase which intracellularly stained with 0.25 ug Coralite@594 Phospho-Mek1 (Thr386) Recombinant Antibody (CL594-81304, Clone:6K5)(red), or 0.25 ug Coralite@594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.