For Research Use Only

## CoraLite®594 Anti-Human CD81 Rabbit Recombinant Antibody

Catalog Number: CL594-98202



**Basic Information** 

Catalog Number: CL594-98202

Concentration: GeneID (NCBI): 100tests, 5 ul/test 975

Source: ENSEMBL Gene ID:
Rabbit ENSG0000110651
Isotype: UNIPROT ID:

IgG P60033
Immunogen Catalog Number: Full Name:

EG1036 CD81 molecule

Calculated MW: 26 kDa

NM\_004356.4

GenBank Accession Number:

Purification Method:

Protein A purification CloneNo.:

241722G4

Recommended Dilutions: FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Applications** 

**Tested Applications:** 

FC

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

## **Background Information**

CD81 (also known as TAPA1 or TSPAN28) is a membrane protein of the tetraspanin superfamily, which are characterized by the presence of four conserved transmembrane regions. Many of these members are expressed on leukocytes and have been implicated in signal transduction, cell-cell interactions, and cellular activation and development. CD81 is involved in signal transduction and cell adhesion in the immune system (PMID: 9597125). CD81 has also been identified as en essential receptor for HCV (hepatitis C virus) (PMID: 21428934).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with 5 ul Coralite®594 Anti-Human CD81 Rabbit RecAb (CL594-98202, Clone: 241722G4) or Coralite®594 Rabbit 1gG Isotype Control RecAb (CL594-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite®594 Anti-Human CD81 Rabbit RecAb (CL594-98202, Clone: 241722G4) (red) or CoraLite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.