## For Research Use Only

## CoraLite®594 Anti-Human KIR2DL3/CD158b2 Rabbit Recombinant Antibody




Catalog Number: CL594-98373-3


Basic Information	Catalog Number: CL594-98373-3	GenBank Accession Number: NM_015868.2	Purification Method: Protein A purification			
	Concentration: 100tests, 5 ul/test	GenelD (NCBI): 3804	CloneNo.: 242282E11			
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: EG2699	UNIPROT ID: P43628-1 Full Name: killer cell immunoglobulin-like receptor, two domains, long cytoplasmic tail, 3	Excitation/Emission maxima wavelengths: 588 nm / 604 nm			
				Calculated MW: 38 kDa		
				Applications	Tested Applications: FC	
		Species Specificity: human				
Background Information	Killer cell immunoglobulin-like receptors (KIRs) are a diverse family of inhibitory and activating receptors expressed on NK cells and a subset of T cells (PMID: 11861603). These polymorphic receptors interact with specific motifs on HLA class I molecules, modulate NK cytolytic activity and are encoded by genes located on chromosome 19q13.4 (PMID: 17069649). KIR2DL3, also known as CD158B2 or NKAT2, is an inhibitory receptor that is specific for HLA-C alleles (HLA-Cw1, HLA-Cw3 and HLA-Cw7) (PMID: 10196125). It is a 341-amino acid transmembrane glycoprotein consisting of an extracellular region containing two C2-type Ig-like domains, a 19-amino acid hydrophobic transmembrane region, and a long cytoplasmic tail with two immunoreceptor tyrosine-based inhibitory motifs (ITIMs). KIR2DL3 inhibits the cytolytic activity of NK cells thus preventing cell lysis (PMID: 7724594).					
Storage	Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide and 0.5% BSA, pH7.3					

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





1x10<sup>^6</sup> human PBMCs were surface stained with CoraLite® Plus 488 Anti-Human NCAM1/CD56 Rabbit Recombinant Antibody (CL488-98033, Clone: 240078A7), and 5 ul CoraLite®594 Anti-Human KIR2DL3/CD158b2 Rabbit RecAb (CL594-98373-3, Clone: 242282E11) or CoraLite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9). Cells were not fixed. Lymphocyte were gated.

1x10^6 human PBMCs were surface stained with 5 ul Coralite®594 Anti-Human KIR2DL3/CD158b2 Rabbit RecAb (CL594-98373-3, Clone: 242282E11) (red) or Coralite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9) (blue). Cells were not fixed. Lymphocyte were gated.