For Research Use Only

CoraLite® Plus 647-conjugated GRP75 Polyclonal antibody

Catalog Number:CL647-14887

Featured Product




Basic Information	Catalog Number: CL647-14887	GenBank Accession Number: BC 000478	Purification Method: Antigen affinity purification	
	Size: 1000 µg/ml	GenelD (NCBI): 3313	Recommended Dilutions: IF/ICC 1:50-1:500	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG6673	UNIPROT ID: P38646 Full Name: heat shock 70kDa protein 9 (morta	Excitation/Emission maxima wavelengths: 654 nm / 674 nm talin)	
				Calculated MW: 74 kDa
		Observed MW: 74-75 kDa		
		Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat	Positive Controls: IF/ICC : HepG2 cells,
Background Informatio	GRP75 (also known as mortalin, HSPA9 or mt-Hsp70) is a constitutively expressed member of the HSPA (HSP70) family of heat-shock proteins. It is located in the mitochondrial matrix and anti-GRP75 is commonly used as the marker for mitochondria. It has been reported that GRP75 is enriched in cancer cells and contributes to carcinogenesis. In addition, decreased expression level of GRP75 has been found in neurodegenerative disorders like Parkinson's disease. This antibody recognized the endogenous GRP75 protein in NIH/3T3 cells. (PMID: 21640711, 22920904)			
Storage	Storage: Store at -20°C. Avoid exposure to Storage Buffer: PBS with 50% Glycerol, 0.05% Pro Aliquoting is unnecessary for -20°		ent.	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 647 GRP75 antibody (CL647-14887) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human GRP75 (CL647-14887) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).