CoraLite® Plus 647-conjugated Cytokeratin 19 Polyclonal antibody





Catalog Number:CL647-14965 Featured Product

Basic Information	Catalog Number: CL647-14965	GenBank Accession Number: BC002539	Purification Method: Antigen affinity purification
	Size: 1000 ug/ml	GenelD (NCBI): 3880	Recommended Dilutions: IF/ICC 1:50-1:500
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG6830	UNIPROT ID: P08727 Full Name: keratin 19	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
		Observed MW: 40 kDa, 46 kDa	
		Applications	Tested Applications:
IF/ICC, FC (Intra) IF/ICC : HeLa of Species Specificity: human, mouse, rat			eLa cells,
Background Information	Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT19, one of type I keratins, is specifically expressed in the periderm, the transiently superficial layer that envelopes the developing epidermis. Due to its high sensitivity, KRT19 is the most used marker for detection of tumor cells disseminated in lymph nodes, peripheral blood, and bone marrow of breast cancer patients. This antibody, generated against full length KRT19 protein, also recognizes KRT17, another type I keratin homologous to KRT19.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human Cytokeratin 19 (CL647-14965) (red), or 0.2 ug CL647-30000 Rabbit IgG (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 647 Cytokeratin 19 antibody (CL647-14965) at dilution of 1:200.