For Research Use Only


CoraLite® Plus 647-conjugated Transgelin-2-specific Polyclonal antibody




Catalog Number:CL647-15508 Featured Product			
Basic Information	Catalog Number: CL647-15508	GenBank Accession Number: NM_003564	Purification Method: Antigen affinity purification
	Size: 1000 μg/ml	Genel D (NCBI): 8407	Recommended Dilutions: IF/ICC 1:50-1:500
	Source: Rabbit	UNIPROT ID: P37802	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
	Isotype: IgG	Full Name: transgelin 2	
		Calculated MW: 22 kDa	
		Observed MW: 22 kDa	
Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human	Positive Controls: IF/ICC : HepG2 cells,	
Background Information	The transgelin family is a group of proteins that belong to 22 kDa actin-related corpnin superfamily. Of all three isoforms, transgelin 1 is the best characterized. Transgelin 1, also known as SM22 alpha, is a specific marker for differentiated smooth muscle cells. Transgelin 2, also known as SM22 beta, is expressed by both smooth muscle and non-smooth muscle cells in a temporally and spatially regulated pattern. Transgelin 3, also known as NP25, is only found in highly differentiated neuronal cells. This antibody can bind transgelin 2 specifically, but not trangelin 1 or 3.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10^6 HepG2 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human Transgelin-2-specific (CL647-15508) (red), or 0.2 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 647 Transgelin-2-specific antibody (CL647-15508) at dilution of 1:200.