For Research Use Only

## CoraLite® Plus 647-conjugated PKC Alpha Polyclonal antibody



Catalog Number: CL647-21991

Featured Product

**Basic Information** 

Catalog Number: CL647-21991

Source: Rabbit

Isotype: IgG

Immunogen Catalog Number:

AG17275

GenBank Accession Number:

AK055431 GenelD (NCBI): 5578

UNIPROT ID: P17252 Full Name:

protein kinase C, alpha

Calculated MW: 77 kDa Observed MW: 77 kDa Purification Method:

Antigen affinity purification
Recommended Dilutions:

IF/ICC: 1:50-1:500

FC (Intra): 0.80 ug per 10<sup>6</sup> cells in a

100  $\mu l$  suspension

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat Positive Controls:

IF/ICC : HepG2 cells,
FC (Intra) : HeLa cells,

## **Background Information**

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. PRKCA is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes.

Storage


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 647 PKC Alpha antibody (CL647-21991) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 647 Anti-Human PKC Alpha (CL647-21991) (red), or 0.8 ug CoraLite® Plus 647-conjugated Rabbit IgG control Rabbit PolyAb (CL647-30000, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.