For Research Use Only

CoraLite® Plus 647-conjugated Beta Catenin Polyclonal antibody



Purification Method:

IF/ICC: 1:50-1:500

100 µl suspension

wavelengths:

654 nm / 674 nm

Antigen affinity purification

Excitation/Emission maxima

FC (Intra): 0.40 ug per 10⁶ cells in a

Recommended Dilutions:

Catalog Number: CL647-51067

Featured Product

1 Publications

Basic Information

Catalog Number: CL647-51067

Source:

Rabbit Isotype:

GenBank Accession Number:

1499

Full Name:

Calculated MW: 781 aa, 86 kDa Observed MW: 92 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

Cited Applications:

Species Specificity: human, mouse, rat, pig

BC058926 GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000168036

UNIPROT ID: P35222

catenin (cadherin-associated protein), beta 1, 88kDa

Positive Controls:

IF/ICC: MCF-7 cells,

FC (Intra): MCF-7 cells, HeLa cells

Background Information

 β -Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. β -Catenin is a 92-kDa protein, originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to a-catenin and the actin cytoskeleton. Besides its essential role in the AJs, $\,\beta$ catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/ β -catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222)

Notable Publications

Author	Pubmed ID	Journal	Application
Xiao-Yan Huang	40531618	Cell Rep	IF

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CoraLite® Plus 647 Beta Catenin antibody (CL647-51067) at dilution of 1:200.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Coralite® Plus 647 Anti-Human Beta Catenin (CL647-51067) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.