For Research Use Only

CoraLite® Plus 647-conjugated Cytokeratin 19 Monoclonal antibody



Catalog Number: CL647-60187

Featured Product

Basic Information

Catalog Number: CL647-60187

Size: 1000 µg/ml Source: Mouse Isotype:

Immunogen Catalog Number:

lgG2b

AG7407

44 kDa Observed MW:

48 kDa

BC002539

3880

P08727

GeneID (NCBI):

UNIPROT ID:

Full Name:

keratin 19

Calculated MW:

GenBank Accession Number:

Purification Method:

Protein A purification

CloneNo.: 3G1E4

Recommended Dilutions:

IF-P 1:50-1:500

Excitation/Emission maxima wavelengths:

654 nm / 674 nm

Applications

Tested Applications: IF-P, FC (Intra) Species Specificity:

human

Positive Controls:

IF-P: human breast cancer tissue, human oesophagus

cancer tissue

Background Information

The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells. KRT19, one of type I keratins, is specifically expressed in the periderm, the transiently superficial layer that envelopes the developing epidermis. Due to its high sensitivity, KRT19 is the most used marker for detection of tumor cells disseminated in lymph nodes, peripheral blood, and bone marrow of breast cancer patients.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed human oesophagus cancer tissue using CoraLite® Plus 647 Cytokeratin 19 antibody (CL647-60187, Clone: 3G1E4) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using Coralite® Plus 647 Cytokeratin 19 antibody (CL647-60187, Clone: 3G1E4) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using Coralite® Plus 647 Cytokeratin 19 antibody (CL647-60187, Clone: 3G1E4) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 647 Anti-Human Cytokeratin 19 (CL647-60187, Clone:3G1E4) (red), or 0.8 ug Mouse IgG2b lostype Control (MPC-11) (65128-1-Ig, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).