For Research Use Only

## CoraLite® Plus 647-conjugated Galectin-3 Monoclonal antibody



Catalog Number: CL647-60207

**Featured Product** 

**Basic Information** 

**Applications** 

Catalog Number: CL647-60207

1000 µg/ml Source: Mouse

lgG2b Immunogen Catalog Number:

AG6753

Isotype:

Calculated MW: 26 kDa Observed MW:

**Tested Applications:** IF/ICC, FC (Intra) Species Specificity:

human, mouse, rat

GenBank Accession Number:

BC001120 Protein A purification GeneID (NCBI): 3958

**UNIPROT ID:** Recommended Dilutions: P17931 IF/ICC 1:50-1:500 Excitation/Emission maxima Full Name:

lectin, galactoside-binding, soluble, 3 wavelengths: 654 nm / 674 nm

CloneNo.:

1C1B2

**Purification Method:** 

Positive Controls: IF/ICC: HeLa cells,

## **Background Information**

Galectins are a family of animal lectins defined by shared characteristic amino-acid sequences and affinity for  $\,\beta$  galactose-containing oligosac-charides (PMID: 8063692). Galectin-3, a member of the  $\beta$ -galactoside-binding proteins, contains one carbohydrate recognition domain (CRD) and a proline- and glycine-rich N-terminal domain through which is able to form oligomers (PMID: 14758078). Galectin-3 is widely expressed in many normal tissues and a variety of tumors. It is found intracellularly in nucleus and cytoplasm or secreted outside of cell, being present on the cell surface or in the extracellular space (PMID: 16478649). Galectin-3 is involved in various biological processes including cell growth, adhesion, differentiation, apoptosis, angiogenesis, immune response, neoplastic transformation and metastasis (PMID: 16478649; 14758078).

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer

31 kDa


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 647 Galectin-3 antibody (CL647-60207, Clone: 1C1B2) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 647 Anti-Human Galectin-3 (CL647-60207, Clone:1C1B2) (red), or 0.8 ug CoraLite® Plus 647 Mouse IgG2b Isotype Control (MPC-11) (CL647-65128, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).