For Research Use Only

CoraLite® Plus 647 Anti-Human CD8a (OKT8)



Catalog Number: CL647-65135

Basic Information

Catalog Number:

CL647-65135

Size:

100tests, 5 $\,\mu$ l/test

Source: Mouse Isotype:

IgG2a, kappa

GenBank Accession Number: BC025715

GeneID (NCBI):

925

ENSEMBL Gene ID: ENSG00000153563

CD8a molecule
Calculated MW:

Full Name:

235 aa, 26 kDa

Purification Method: Affinity purification

CloneNo.: OKT8

Excitation/Emission maxima wavelengths:

654 nm / 674 nm

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA, pH 7.3.

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with PacificBlue Anti-Human CD3 and 5 ul Coralite® Plus 647 Anti-Human CD8a (CL647-65135, Clone: OKT8), or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 647 Anti-Human CD8a (CL647-65135, Clone: OKT8), or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.