For Research Use Only

CoraLite® Plus 647-conjugated HDAC1 Monoclonal antibody



Catalog Number:CL647-66085 1 Publications

Basic Information

Catalog Number:

GenBank Accession Number: BC000301

Purification Method:

CL647-66085

GeneID (NCBI):

Protein G purification

Size:

CloneNo.:

1000 µg/ml

3065

4C4G7

Source: Mouse

UNIPROT ID: Q13547

Recommended Dilutions:

Isotype:

Full Name:

IF/ICC 1:500-1:2000

lgG1

histone deacetylase 1

Excitation/Emission maxima

Immunogen Catalog Number:

Calculated MW:

wavelengths: 654 nm / 674 nm

AG19128

55 kDa

Observed MW:

65 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

Positive Controls:

IF/ICC: A431 cells,

Species Specificity: human, mouse, rat

Cited Species:

mouse

Background Information

Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex, which is responsible for gene expression silencing. It also plays an important role in the control of cell proliferation and differentiation by interacting with RB, p53 and other transcription factors. At least 4 classes of HDAC were identified. As a class I HDAC, HDAC 1 was primarily found in the nucleus. This antibody is raised against residues near the C terminus of human HDAC1. The calculated molecular weight of HDAC1 is 55 kDa, but modified HDAC1 is about 60-65 kDa.

Notable Publications

Author Application **Pubmed ID** Journal J Adv Res Chun Zhang 39107200


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite® Plus 647 HDAC1 antibody (CL647-66085, Clone: 4C4G7) at dilution of 1:1000.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 647 Anti-Human HDAC1 (CL647-66085, Clone:4C4G7) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).