For Research Use Only

CoraLite® Plus 647-conjugated SURVIVIN Monoclonal antibody


Catalog Number:CL647-66495 Featured Product



Basic Information	Catalog Number: CL647-66495	GenBank Accession Number: BC008718	Purification Method: Protein A purification
	Size:	GenelD (NCBI):	CloneNo.:
	1000 µg/ml	332	3B9H7
	Source: Mouse	UNIPROT ID: O15392	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
	Isotype: IgG1 Immunogen Catalog Number: AG20958	Full Name: baculoviral IAP repeat-containing 5 Calculated MW: 16 kDa	
		Applications	
Species Specificity: Human			
Background Information	Survivin, also called BIRC5, is a unique member of the inhibitor of apoptosis (IAP) protein family. Survivin is a 16 kDa anti-apoptotic protein highly expressed during fetal development and cancer cell malignancy, but is completely absent in terminally differentiated cells. The differential expression of survivin in cancer versus normal tissues makes it a useful tool in cancer diagnosis and a promising therapeutic target. Survivin expression is also highly regulated by the cell cycle and is only expressed in the G2-M phase. It is known that survivin localizes to the mitotic spindle by interaction with tubulin during mitosis and may play a contributing role in regulating mitosis. Disruption of survivin-microtubule interactions results in loss of survivin's anti-apoptosis function and increased caspase-3 activity, a mechanism involved in cell death, during mitosis. It also is a direct target gene of the Wnt pathway and is upregulated by beta-catenin.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human SURVIVIN (CL647-66495, Clone:3B9H7) (red), or 0.2 ug Control Antibody. (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).