CoraLite® Plus 647-conjugated FBL Monoclonal antibody

Antibodies | ELISA kits | Proteins www.ptglab.com


Catalog Number: CL647-66985

| Basic Information | Catalog Number: CL647-66985 | GenBank Accession Number: BC019260 | Purification Method: Protein G purification |
|-----------------------|---|--|--|
| | Size: 1000 µg/ml | GenelD (NCBI): 2091 | CloneNo.: 3A9E8 |
| | Source: Mouse | UNIPROT ID: P22087 | Recommended Dilutions: IF/ICC 1:50-1:500 |
| | Isotype: IgG1 Immunogen Catalog Number: AG8841 | Full Name: fibrillarin | Excitation/Emission maxima wavelengths: 654 nm / 674 nm |
| | | Calculated MW: 321 aa, 34 kDa | |
| | | Observed MW: 34 kDa | |
| Applications | Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat | Positive Controls: | |
| | | IF/ICC : H | epG2 cells, |
| | Fibrillarin (Fbl) is an essential nucleolar protein that plays an essential role in ribosome biogenesis and more particularly in the methylation of ribosomal RNAs and rDNA histones. FBL is highly expressed in various cancers and correlates with poor survival outcomes in cancer patients, such as hepatocellular carcinoma(PMID: 33376525). Knockdown of FBL sensitizes tumor cells and xenografts to DNA crosslinking agents, and leads to homologous recombination-mediated DNA repair defects(PMID: 37489617). | | |
| Background Informatio | correlates with poor survival outco Knockdown of FBL sensitizes tumo | ribosomal RNAs and rDNA histones. F omes in cancer patients, such as hepat or cells and xenografts to DNA crosslin | BL is highly expressed in various cancers and cocllular carcinoma(PMID: 33376525). |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 647 FBL antibody (CL647-66985, Clone: 3A9E8) at dilution of 1:200.



1x10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 647 Anti-Human FBL (CL647-66985, Clone:3A9E8)(red), or 0.4 ug CoraLite® Plus 647 Mouse IgG1 Isotype Control (MOPC-21) (CL647-65124, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).