For Research Use Only

CoraLite® Plus 647 Anti-Human CD97 Rabbit Recombinant Antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: CL647-98094-2

Basic Information

Catalog Number: CL647-98094-2

Concentration: 100tests, 5 ul/test

Source: Rabbit

Isotype: IgG GenBank Accession Number: BC026690

GeneID (NCBI):

UNIPROT ID: P48960

Full Name: CD97 molecule

Calculated MW: 722 aa, 79 kDa Purification Method:

Protein A purification

CloneNo.: 241335D8

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications:

. .

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

Background Information

CD97 is a member of the seven-trans-membrane subfamily of the class B G protein-coupled receptor (GPCR) group of the epidermal growth factor (EGF) and is present on the surface of lymphocytes, monocytes, macrophages, dendritic cells, granulocytes, and smooth muscle, being a dimeric glycoprotein with a 75-90 kDa intracellular domain and a 28 kDa extracellular domain (PMID: 23969601). It plays a vital role in cell adhesion, migration, and cell connection regulation (PMID: 36464675). CD97 is expressed on T-cells, but rarely on B cells. CD97 might play key roles in cellular adhesion, through connections with other proteins of the cell surface and of the extracellular matrix (PMID: 32781778).

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data





1x10^6 human PBMCs were surface stained with CoraLite® Plus 488 Anti-Human CD3 (UCHT1) Mouse IgG2a Recombinant Antibody (CL488-65570, Clone: UCHT1), and 5 ul CoraLite® Plus 647 Anti-Human CD97 Rabbit RecAb (CL647-98094-2, Clone: 241335D8) or CoraLite® Plus 647 Rabbit IgG Isotype Control RecAb (CL647-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul Coralite® Plus 647 Anti-Human CD97 Rabbit RecAb (CL647-98094-2, Clone: 241335D8) (red) or Coralite® Plus 647 Rabbit IgG Isotype Control RecAb (CL647-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.