For Research Use Only

## CoraLite® Plus 647 Anti-Human IFN-gamma Rabbit Recombinant Antibody



Catalog Number: CL647-98187

**Basic Information** 

Catalog Number: GenBank Accession Number: CL647-98187 BC070256

Concentration: GeneID (NCBI): 100tests, 5 ul/test 3458

Source: ENSEMBL Gene ID: Rabbit ENSG00000111537

Isotype: UNIPROT ID: IgG P01579

Immunogen Catalog Number:Full Name:EG0004IFN gamma

Calculated MW: 166 aa, 19 kDa Purification Method:

Protein A purification CloneNo.:

241390B10

Recommended Dilutions:

FC (Intra): 5 ul per 10^6 cells in a 100  $\,$ 

µl suspension

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

Tested Applications:

FC (Intra)

Species Specificity:

human

Positive Controls:

FC (Intra): PMA and ionomycin treated human PBMCs,

## **Background Information**

Interferon gamma (IFN  $\gamma$  ), is a type II interferon that provides immunity against bacterial, viral and protozoan infections. In its active form, IFN  $\gamma$  is a glycosylated, non-covalently linked homodimer of 29-32 kDa subunits. It is produced by a number of immune cell types including natural killer cells, natural killer T cells, and effector lymphocyte T cells following antigenic and inflammatory triggers. The IFN  $\gamma$  dimer binds to its cognate receptor which has two subunits: IFN- $\gamma$  R1 which is the ligand-binding chain ( $\alpha$  chain) and IFN- $\gamma$  R2, the signal-transducing chain ( $\beta$  chain). Binding to the receptor activates the JAK/STAT pathway which in turn activates IFN  $\gamma$  responsive genes. While IFN  $\gamma$  can inhibit viral replication, it also works as an immune-modulator and immune-stimulator by increasing surface expression of class I MHC proteins. (PMID: 19268625; 10688427)


Storage

Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## Selected Validation Data



1x10^6 untreated or PMA and ionomycin treated human PBMCs were intracellularly stained with 5ul Coralite® Plus 647 Anti-Human IFN-gamma Rabbit RecAb (CL647-98187, Clone: 241390B10) . Cells were co-stained with APC Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (APC-65569, Clone: OKT3). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.