For Research Use Only

CoraLite®700 Anti-Human CD7 Rabbit Recombinant Antibody



Catalog Number: CL700-98204

Basic Information

Catalog Number: CL700-98204

Concentration: 100tests, 5 ul/test

Source: Rabbit

Isotype: IgG

Immunogen Catalog Number:

EG3034

GenBank Accession Number:

NM_006137.7 GeneID (NCBI):

924

ENSG00000173762

UNIPROT ID: P09564 Full Name:

CD7 molecule

Calculated MW: 25 kDa

Purification Method:

Protein A purification

CloneNo.: 242402H8

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 μ l

suspension

Excitation/Emission maxima

wavelengths: 702 nm / 723 nm

Applications

Tested Applications:

FC

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

Background Information

CD7, also known as Leu-9 or GP40, is a 40-kDa single-pass type I transmembrane glycoprotein belonging to the immunoglobulin superfamily. CD7 is expressed on thymocytes, T cells, NK cells, and cells in the early stages of T, B, and myeloid cell differentiation (PMID: 10530432). It is one of the earliest antigens to appear on cells of the T-lymphocyte lineage (PMID: 3501369). CD7 plays a significant role in T-cell and T-cell/B-cell interactions during early lymphoid development. It is involved in the regulation of cell signaling and biology, including T-cell activation, proliferation, and the expression of interleukin-2 receptor alpha (IL-2R a) (PMID: 11485208). CD7 is not only a marker for T-cells but also has functional implications in immune cell interactions and adhesion. It can provide co-stimulatory signals and is associated with CD3 and CD45, participating in T-cell activation (PMID: 7523512).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3 $\,$

Selected Validation Data



1x10^6 human PBMCs were surface stained with CoraLite® Plus 488 Anti-Human CD4, and 5 ul CoraLite® 700 Anti-Human CD7 Rabbit RecAb (CL700-98204, Clone: 242402H8) or unstained. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite®700 Anti-Human CD7 Rabbit RecAb (CL700-98204, Clone: 242402H8) (red) or unstained (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.