For Research Use Only

CoraLite® Plus 750-conjugated DYKDDDDK tag Polyclonal antibody (Binds to FLAG® tag epitope)




Catalog Number:CL750-20543			
Basic Information	Catalog Number: CL750-20543	GenBank Accession Number:	Purification Method: Antigen affinity purification
	Size: 1000 µg/ml	GeneID (NCBI): Full Name:	Recommended Dilutions: WB 1:2000-1:16000
	Source: Rabbit		Excitation/Emission maxima wavelengths: 755 nm / 780 nm
	lsotype: IgG		
	Immunogen Catalog Number: AG2329		
Applications	Tested Applications:	Positive Co	ntrols:
	WB Species Specificity: recombinant protein	WB : Transfected HEK-293 cells,	
Background Information	Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. The DYKDDDDK(FLAG) peptide has been used extensively as a general tag in expression vectors. This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion. N-terminal DDDDK vectors provide an Ek cleavage site for removal of the fusion tag. The DDDDK peptide is likely to be located on the surface of a fusion protein because of its hydrophilic nature. As a result, the DDDDK peptide is more likely to be accessible to antibodies. A DDDDK-tag can be used in many different assays that require recognition by an antibody, such as western blotting, immunocytochemistry, immunoprecipitation, flow cytometry, protein purification, and in the study of protein-protein interactions, cell ultrastructure, and protein localization and so on. This antibody is a rabbit polyclonal antibody raised against 3xFlag (3xDYKDDDDKT) sequence and recognizes the (1x) and (3x)DYKDDDDK peptide and detects DDDDK-tagged proteins. Anti-FLAG is a registered trademark of Sigma-Aldrich Biotechnology.		
Storage	Storage: Store at -20°C. Avoid exposure to lig Storage Buffer: PBS with 50% Glycerol, 0.05% Procl Aliquoting is unnecessary for -20°C	•	<u>.</u>

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-20543 (DYKDDDDK tag antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.