For Research Use Only

CoraLite® Plus 750-conjugated PCNA Monoclonal antibody


Catalog Number:CL750-60097 Featured Product



Basic Information	Catalog Number: CL750-60097	GenBank Accession Number: BC000491	Purification Method: Protein G purification
	Size: 1000 ug/ml	GenelD (NCBI): 5111	CloneNo.: 10D10E11
	Source: Mouse	UNIPROT ID: P12004	Recommended Dilutions: WB 1:2000-1:16000
	Isotype: IgG1 Immunogen Catalog Number: AG7416	Full Name: proliferating cell nuclear antigen Calculated MW: 29 kDa/31 kDa	Excitation/Emission maxima wavelengths: 755 nm / 780 nm
		Applications	
Background Information	Proliferating Cell Nuclear Antigen, commonly known as PCNA, is a protein that acts as a processivity factor for DNA polymerase δ in eukaryotic cells. This protein is an auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. PCNA induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. It has to be loaded onto DNA in order to be able to stimulate APEX2. PCNA protein is highly conserved during evolution; the deduced amino acid sequences of rat and human differ by only 4 of 261 amino acids. PCNA has been used as loading control for proliferating cells. The calculated molecular weight of PCNA is 29 kDa, but modified PCNA is 36kDa (PMID: 1358458).		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-60097 (PCNA antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.