#### For Research Use Only

# CoraLite® Plus 750 Anti-Mouse CD8a (53-6.7)



Catalog Number: CL750-65069

1 Publications

**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 CL750-65069
 BC030679

 Concentration:
 GeneID (NCBI):

 100ug, 0.5 mg/ml
 12525

 Source:
 UNIPROT ID:

 Post 774

Rat P01731
Isotype: Full Name:

IgG2a, kappa CD8 antigen, alpha chain

Purification Method:

Affinity purification CloneNo.:

53-6.7

Recommended Dilutions: FC: 0.5 ug per 10^6 cells in 100  $\,\mu$  l

suspension

Excitation/Emission maxima

wavelengths: 755 nm / 780 nm

**Applications** 

**Tested Applications:** 

FC

Cited Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC: mouse splenocytes,

### **Background Information**

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer, while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Pu Wang	39859511	Int J Mol Sci	FC

Storage


Storage:

Store at 2-8  $^{\circ}\text{C}$  . Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 0.09% sodium azide, pH7.3

## Selected Validation Data



1X10^6 mouse splenocytes were surface stained with CoraLite® 488 Plus Anti-Mouse CD4 (GK1.5) (CL488-65104, Clone: GK1.5) and 0.5 ug CoraLite® Plus 750 Anti-Mouse CD8a (CL750-65069, Clone: 53-6.7) or 0.5 ug isotype control. Cells were not fixed.