For Research Use Only

## CoraLite® Plus 750 Anti-Mouse CD8a (53-6.7)



Catalog Number: CL750-65069

**Basic Information** 

Catalog Number:

GenBank Accession Number: BC030679

Purification Method:

CL750-65069

DC0300/9

Affinity purification

Size:

GeneID (NCBI): 12525 CloneNo.: 53-6.7

100ug, 0.5 mg/ml Source:

UNIPROT ID:

Excitation/Emission maxima

Isotype:

P01731 Full Name: wavelengths: 755 nm / 780 nm

IgG2a, kappa

CD8 antigen, alpha chain

**Applications** 

**Tested Applications:** 

FC

Species Specificity:

Mouse

**Background Information** 

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide.

## Selected Validation Data



1X10^6 mouse splenocytes were surface stained with CoraLite® 488 Plus Anti-Mouse CD4 (GK1.5) (CL488-65104, Clone: GK1.5) and 0.5 ug CoraLite® Plus 750 Anti-Mouse CD8a (CL750-65069, Clone: 53-6.7) or 0.5 ug isotype control. Cells were not fixed.