For Research Use Only

CoraLite® Plus 750 Anti-Human CD73 (AD2) Mouse IgG2a Recombinant Antibody



Catalog Number: CL750-65564

Basic Information

Catalog Number: CL750-65564

Size:

100tests, 5 ul/test

Source: Mouse Isotype:

lgG2a

GenBank Accession Number: BC015940

GeneID (NCBI):

4907

UNIPROT ID: P21589 Full Name:

5'-nucleotidase, ecto (CD73)

Calculated MW: 29 kDa, 63 kDa Purification Method:

Protein A purification

CloneNo.: AD2

Excitation/Emission maxima

wavelengths: 755 nm / 780 nm

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane-bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 human PBMCs were surface stained with PE Anti-Human CD45,and 5 ul Coralite® Plus 750 Anti-Human CD73 (AD2) Mouse IgG2a Recombinant Antibody (CL750-65564, Clone: AD2) or Coralite® Plus 750 Mouse IgG2a Isotype Control (C1.18.4) (CL750-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 750 Anti-Human CD73 (AD2) Mouse IgG2a Recombinant Antibody (CL750-65564, Clone: AD2) (red) or CoraLite® Plus 750 Mouse IgG2a Isotype Control (C1.18.4) (CL750-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not