CoraLite® Plus 750-conjugated Beta Actin Monoclonal antibody



Catalog Number:CL750-66009


Featured Product


Basic Information	Catalog Number: CL750-66009	GenBank Accession Number: NM_001101	Purification Method: Protein A purification
	Size: 1000 µg/ml	GeneID (NCBI): 60	CloneNo.: 2D4H5
	Source: Mouse	UNIPROT ID: P60709	Recommended Dilutions: WB 1:500-1:2000
	Isotype: IgG2b	Full Name: actin, beta	Excitation/Emission maxima wavelengths: 755 nm / 780 nm
		Calculated MW: 42 kDa	
		Observed MW: 42 kDa	
Applications	Tested Applications: WB Species Specificity: human, mouse, rat	Positive Controls:	
		WB : mouse kidney tissue, mouse brain tissue, HeLa cells, NIH/3T3 cells, HSC-T6 cells, Jurkat cells	
	Actins are highly conserved globular proteins that are involved in various types of cell motility and are ubiquitousl expressed in all eukaryotic cells. At least six isoforms of actins are known in mammals and other vertebrates: alph (ACTC 1, cardiac muscle 1), alpha 1 (ACTA1, skeletal muscle) and 2 (ACTA2, aortic smooth muscle), beta (ACTB), gamma 1 (ACTG1) and 2 (ACTG2, enteric smooth muscle). Beta and gamma 1 are two non-muscle actin proteins. Most actins consist of 376aa, while ACTG2 (rich in muscles) has 375aa and ACTG1(found in non-muscle cells) has only 374aa. Beta actin has been widely used as the internal control in RT-PCR and Western Blotting as a 42-kDa protein. However, the 41 kDa cleaved fragment of beta actin can be generated during apoptosis process. This antibody can recognize all the actins. The isotype of this antibody is IgG2b. For murine tissue sample, conjugated antibody (HRP-66009) or rabbit antibody (20536-1-AP) is preferable.		
Background Information	(ACTC1, cardiac muscle 1), al gamma 1 (ACTG1) and 2 (AC Most actins consist of 376aa, only 374aa. Beta actin has be protein. However, the 41 kDa antibody can recognize all th	TG2, enteric smooth muscle). Beta and gau while ACTG2 (rich in muscles) has 375aa en widely used as the internal control in I cleaved fragment of beta actin can be ge e actins. The isotype of this antibody is Ig	TA2, aortic smooth muscle), beta (ACTB), mma 1 are two non-muscle actin proteins. and ACTG1(found in non-muscle cells) has RT-PCR and Western Blotting as a 42-kDa merated during apoptosis process.This

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with CL750-66009 (Beta Actin antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Various lysates were subjected to SDS PAGE followed by western blot with CL750-66009 (Beta Actin antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.