## For Research Use Only

## CoraLite® Plus 750-conjugated Vinculin Monoclonal antibody

Catalog Number: CL750-66305




Basic Information	Catalog Number: CL750-66305	GenBank Accession Number: BC039174	Purification Method: Protein G purification				
	Size: 1000 ug/ml	GenelD (NCBI): 7414	CloneNo.: 2B5A7				
	Source: Mouse	UNIPROT ID: P18206	Recommended Dilutions: WB 1:500-1:2000				
	Isotype: IgG1 Immunogen Catalog Number: AG24946	Full Name: vinculin Calculated MW: 1133 aa, 124 kDa Observed MW: 117 kDa	Excitation/Emission maxima wavelengths: 755 nm / 780 nm				
				Applications	Tested Applications: WB	Positive Controls:	
					Species Specificity: human, mouse, rat, pig	WB : K-562 cells, PC-3 cells	
Background Informati	involved in cell-matrix adhesion potentiates mechanosensing by th locomotion. Vinculin is a 117-kDa	Vinculin belongs to the vinculin/alpha-catenin family. It is an actin filament (F-actin)-binding protein which involved in cell-matrix adhesion and cell-cell adhesion. Vinculin regulates cell-surface E-cadherin expression and potentiates mechanosensing by the E-cadherin complex. It may also play important roles in cell morphology and locomotion. Vinculin is a 117-kDa, 1,066-amino-acid protein which is ubiquitously expressed. Its splice variant, metavinculin (124 kDa), is muscle-specific.					
	Storage:						
Storage	Store at -20°C. Avoid exposure to Storage Buffer: PBS with 50% Glycerol, 0.05% Pro		nt.				

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-66305 (Vinculin antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.