## For Research Use Only

## CoraLite® Plus 750-conjugated Phospho-JUN (Ser73) Recombinant antibody




Catalog Number:CL750-80086


Basic Information	Catalog Number: CL750-80086	GenBank Accession Number: BC068522	Purification Method: Protein A purification
	Concentration: 1000 ug/ml	GenelD (NCBI): 3725	CloneNo.: 4A18
	Source: Rabbit	UNIPROT ID: P05412	Recommended Dilutions: WB 1:500-1:1000
	Isotype: IgG	Full Name: jun oncogene Calculated MW: 331 aa, 36 kDa	Excitation/Emission maxima wavelengths: 755 nm / 780 nm
		Applications	
WB : SH-SY5Y cells, Species Specificity: human, mouse			
	JUN, the most extensively studied protein of the activator protein-1 (AP-1) complex, is involved in numerous cell activities, such as proliferation, apoptosis, survival, tumorigenesis and tissue morphogenesis (PMID: 22180088). JUN is a transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. It promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation. JUN is a basic leucine zipper (bZIP) transcription factor that acts as homo- or heterodimer, binding to DNA and regulating gene transcription (PMID: 9732876). In additon, extracellular signals can induce post-translational modifications of JUN, resulting in altered transcriptional activity and target gene expression (PMID:8464713). More over, it has uncovered multiple layers of a complex regulatory scheme in which JUN is able to crosstalk, amplify and integrate different signals for tissue development and disease. Jun is predominantly nuclear, ubiquitinated Jun colocalizes with lysosomal proteins (PMID: 15469925). This antibody is raised against synthetic phosphopeptide corresponding to residues surrounding Ser73 of human JUN, which can detect the bands around 42-45 kDa.		
Background Informatior	activities, such as proliferation is a transcription factor that re- activity of NR5A1 when phosp signaling pathway stimulation heterodimer, binding to DNA- can induce post-translational expression (PMID:8464713). N JUN is able to crosstalk, ampl predominantly nuclear, ubiqu raised against synthetic phosp	n, apoptosis, survival, tumorigenesis and cognizes and binds to the enhancer hepta whorylated by HIPK3 leading to increased n. JUN is a basic leucine zipper (bZIP) tran and regulating gene transcription (PMID: 9 modifications of JUN, resulting in altered fore over, it has uncovered multiple layers ify and integrate different signals for tissc itinated Jun colocalizes with lysosomal p phopeptide corresponding to residues surr	tissue morphogenesis (PMID: 22180088). JUN mer motif 5'-TGA[CG]TCA-3'. It promotes steroidogenic gene expression upon cAMP scription factor that acts as homo- or 732876). In additon, extracellular signals transcriptional activity and target gene s of a complex regulatory scheme in which ie development and disease. Jun is roteins (PMID: 15469925). This antibody is

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





1X10^6 NIH/3T3 cells untreated (dashed lines) or treated with UV which intracellularly stained with 0.25 ug CoraLite® Plus 750-conjugated Phospho-JUN (Ser73) Recombinant antibody (CL750-88086, Clone:4A18)(red), or 0.25 ug CoraLite® Plus 750 Rabbit 1gG Isotype Control RecAb (CL750-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.

Various lysates were subjected to SDS PAGE followed by western blot with CL750-80086 (Phospho-JUN (Ser73) antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.